arXiv:1612.00177v1 [math.CV] 1 Dec 2016

One sided conformal collars and the reflection principle

V. Liontou and V. Nestoridis

December 2, 2016

Abstract

If a Jordan curve σ has a one-sided conformal collar with "good" properties, then, using the Reflection principle, we show that any other conformal collar of σ from the same side has the same "good" properties. A particular use of this fact concerns analytic Jordan curves, but in general the Jordan arcs we consider do not have to be analytic. We show that if an one-sided conformal collar bounded by σ is of class A^p, then any other collar bounded by σ and from the same side of σ is of class A^p.

A.M.S classification no: primary 30C99, 30E25 secondary30H10

Key words and phrases: Reflection principle, one sided conformal collar, analytic curves, A^p spaces.

1 Introduction

In this paper, for two conformal mappings Φ and F, we write $F = \Phi \circ (\Phi^{-1} \circ F)$ whenever this has a meaning, as explained later. Using our assumptions, the function $\Phi^{-1} \circ F$ maps a circular arc τ on another circular arc. Thus, by Reflection $h = \Phi^{-1} \circ F$ extends to an injective holomorphic function on a neighbourhood V of a part of τ. Thus, h' extends continuously on τ and does not vanish at any point of τ. Since $F = \Phi \circ h$ and $F' = (\Phi' \circ h)h'$, if Φ or Φ' has some good properties we can transfer them to F and F'. The first application of this fact is described below.
Let J be a Jordan curve in $\mathbb{C}$, that is a homeomorphic image of the unit

circle T defined by $\gamma : T \to J$. The Jordan curve J is called analytic if there is a two sided conformal collar around it . That is, if there exists an annulus $D(0, r_1, r_2) = \{z \in \mathbb{C} : r_1 < |z| < r_2\}, 0 < r_1 < 1 < r_2$ and an injective holomorphic mapping $\phi : D(0, r_1, r_2) \to \mathbb{C}$ such that $\phi(e^{i\theta}) = \gamma(e^{i\theta})$ for all $e^{i\theta} \in T$, $\theta \in \mathbb{R}$. A consequence of the existence of the above two sided conformal collar is that, if Ω is the Jordan region bounded by J and $\Phi : D \to \Omega$ is a Riemann mapping from the open disk D onto Ω, then Φ has a conformal extension on a larger disc $D(0, r) = \{z \in \mathbb{C} : |z| < r\}, r > 1$. In particular, the function Φ extends continuously on the unit circle T and the same holds for the derivative Φ' and $\Phi'(e^{i\theta}) \neq 0$ for all $e^{i\theta} \in T$.
In the present paper we investigate which conclusions still hold if we do not assume the existence of a two-sided conformal collar around the Jordan curve J but we assume the existence of a one sided conformal collar bounded by J.
Thus, we assume the existence of an injective holomorphic function $\phi : D(0, r, 1) \underline{\to \mathbb{C},\ D}(0, r, 1) = \{z \in \mathbb{C} : r < |z| < 1\}$ which extends continuously on $\overline{D(0, r, 1)}$ and let us call ϕ the extension as well. We also assume that ϕ is one to one on $D(0, r, 1) \cup T$ where $T = \{z \in \mathbb{C} : |z| = 1\}$ and that $\phi_{|_T} : T \to J$ is a homeomorphism. Futhermore,we assume that ϕ' has a continuous extension on $\overline{D(0, r, 1)}$ and that $\phi'(e^{i\theta}) \neq 0$ for all $\theta \in \mathbb{R}$. Then we show that the Riemann map $\Phi : D \to \Omega$, onto the Jordan region Ω bounded by J has similar properties; that is, Φ and Φ' extend continuously on $\overline{D}$ and $\Phi'(e^{i\theta}) \neq 0$ for all $\theta \in \mathbb{R}$. Certainly ,the fact that Φ extends continuously on $\overline{D}$ is a consequence of the Ozgood-Caratheodory theorem [6]. The rest is a consequence of the Reflection Principle ([1]). We also can extend the above fact for every finite number $p + 1$ of derivatives, provided that the one-sided conformal collar is of class A^p.([3],[5]) Furthermore, instead of considering the unit circle we can replace it by one-sided free boundary arcs ([1])which are analytic arcs and obtain similar results (see Th.2.2 below)
We also mention that our result relates to the considerations of ([2])

Finally, a second application of the method of the present paper can be found in [7], see proof of Th. 4.2 where it is proven the following. If Ω is a domain in $\mathbb{C}$, whose boundary contains an analytic arc J and the boundary values of a holomorphic function $f \in H(\Omega)$ define a function in $C^p(J)$, then the derivatives $f^{(l)}(z), z \in \Omega, l = 0, ..., p$ extend continuously on J.

2 One sided conformal collar

We start with the following definition.

Definition 2.1. *A complex function $\phi(t)$ of a real variable t defined on an interval $a < t < b$, defines an analytic curve if, for every t_0 in the (a, b),the derivative $\phi'(t_0)$ exists and it is non zero and ϕ has a representation*

$$\phi(t) = \sum_{n=0}^{\infty} \alpha_n(t_0)(t - t_0)^n, \alpha_n(t_0) \in \mathbb{C},$$

where the power series converges in some interval $(t_0-p, t_0+p), p = p(t_0) > 0$, and coincides with ϕ in this interval.

But if this is so, then from Abel's lemma, the series is also convergent for complex values of t, as long as $|t-t_0| < p$ and represents an analytic function on that disk. In overlapping disks the functions are the same, because they coincide on a segment of the real axis. Thus the function $\phi(t)$ can be extended as an analytic and injective function in a neighbourhood of (a, b) in $\mathbb{C}$.[1]

Theorem 2.2. *Let W_1 and W_2 be regions and $J_1$and J_2 be analytic arcs of the boundaries of the regions W_1, W_2 respectively. We also assume that there are neighbourhoods U_1, U_2 of J_1, J_2 respectively for which $U_1 \cap W_1$ and $U_2 \cap W_2$ are connected regions. For the analytic arcs J_1 and J_2 we assume that:*

1. *For every point of J_1 or J_2 there exists a neighbourhood whose intersection with the whole boundary of W_1 or W_2 respectively is the same with its intersection with J_1 or J_2.*

2. *For every point z in $J_i, i = 1, 2$ and for every disk $D(z, \epsilon)$ such that $D(z, \epsilon) \cap \partial W_i = D(z, \epsilon) \cap J_i$, the arc J_i separates $D(z, \epsilon)$ in two subsets, one completely inside W_i and the other completely outside W_i.*

3. *The arcs $J_i, i = 1, 2$ are simple analytic and the injective maps ϕ_1, ϕ_2 with $J_1 := \phi_1 : (a_1, b_1) \to \mathbb{C}$ and $J_2 := \phi_2 : (a_2, b_2) \to \mathbb{C}$ satisfy that ϕ_1' and ϕ_2' exist and are non-zero for every point of the segments (a_1, b_1) and (a_2, b_2) respectively*

We also assume that a function $\phi : W_1 \cup J_1 \to \mathbb{C}$ is injective, continuous on $W_1 \cup J_1$ and holomorphic in W_1 and we denote $G := \phi(W_1)$. We suppose that

ϕ' can be extended in $W_1 \cup J_1$ continuously. Let $\gamma = \phi(J_1)$ and let a function $F : W_2 \to \mathbb{C}$ be continuous on $W_2 \cup J_2$ and holomorphic on W_2. Suppose that for all z in $\phi(J_1)$, there exists a $\epsilon > 0$ so that $D(z, \epsilon) \cap F(W_2) \subseteq G$ and if a sequence $(z_n)_{n\in\mathbb{N}}, z_n \in W_2$ for all $n \in \mathbb{N}$ accumulates only in J_2, then $F(z_n)$ accumulates only in γ and $F(J_2) \subseteq \gamma = \phi(J_1)$.
Then:

(a) The function F', can be continuously extended in $W_2 \cup J_2$.

(b) If $\phi'(z) \neq 0$ for all z in J_1 and F is injective in W_2 then $F'(\zeta) \neq 0$ for all ζ in J_2

Proof. First we prove statement (a).
Let $z \in J_2$. There exists a $\delta_z > 0$ such that $D(z, \delta_z) \cap (\partial W_2 \setminus J_2) = \emptyset$. Therefore, the function

$$h := \phi^{-1} \circ F : D(z, \delta_z) \cap W_2 \to W_1$$

is well defined. The function h is holomorphic, since F is holomorphic in W_2. The function

$$\phi^{-1} : \phi(W_1) \to W_1$$

is holomorphic also holomorphic in $\phi(W_1)$ and if $\delta_z > 0$ is small enough, then $F\big(D(z, \delta_z) \cap W_2\big) \subseteq \phi(W_1)$, for every $z \in \gamma$.
We assume that $\{z_n, n \in \mathbb{N}\}$ is a sequence of points in W_2. We are going to prove that if all the accumulation points of $\{z_n\}_{n\in\mathbb{N}}$ belong in J_2, then all the accumulation points of sequence $b_n := h(z_n), n \in \mathbb{N}$ belong in J_2. From our hypothesis it follows that $F(z_n), n \in \mathbb{N}$ accumulates in γ. It suffices to prove that $\{\phi^{-1} \circ F(z_n), n \in \mathbb{N}\}$ accumulates in J_1. If we suppose that there is an accumulation point w of the sequence $b_n := \phi^{-1} \circ F(z_n)$ that does not belong in J_1, then ,there exists a subsequence $\{b_{k_n}, n \in \mathbb{N}\}$ that converges to $w \in W_1$. Thus, $\phi(b_{k_n})$ defines a subsequence of $F(z_n)$ in $\phi(W_1)$. Since ϕ is continuous on $W_1 \cup J_1$,then $\phi(z_{k_n})$ converges to $\phi(w)$. Since all accumulation points of $F(z_n)$ belong in γ, therefore $\phi(w)$ belongs in γ. Thus, there exists an $z_1 \in J_1$ such as $\phi(z_1) = \phi(w)$. This is impossible since z_1 is on the boundary of W_1 and w inside of W_1 and Φ is injective on $W_1 \cup J_1$.
Therefore if $(z_n)_n$ accumulates on J_2, $h(z_n)$ accumulates on J_1.

Finally, h is holomorphic on $D(z, \delta_z) \cap W_2$ and continuous on $\overline{D(z, \delta_z) \cap W_2}$. According to the Reflection Principle for analytic arcs [1], h is holomorphic

in $D(z,\delta'_z)$for some $\delta'_z, 0 < \delta'_z \leq \delta_z$. Therefore $F = \phi \circ \phi^{-1} \circ F$ is continuous in J_2 since ϕ can be continuously extended in J_1.
Considering F restricted in $D(z,\delta_z) \cap W_2$ it is analytic and

$$F'(z) = [\phi \circ (\phi^{-1} \circ F)]'(z) = \phi' \circ (\phi - 1 \circ F)(z) \cdot (\phi^{-1} \circ F)'(z)$$

$$\forall z \in D(z,\delta_z) \cap W_2$$

Since $\phi\prime$ is continuous in J_1 and from the Reflection Principle $(\phi^{-1} \circ F)'$ is holomorphic in $D(z,\delta_z)$, therefore F' is continuous on $\overline{D(z,\delta_z) \cap W_2}$.

The above statements are valid for all $z \in J_2$. From analytic continuation the extensions of $h \in D(z_i,\delta'_{z_i}) \cap D(z_j,\delta'_{z_j}) \cap W_2$ match, thus they also match in $D(z_i,\delta'_{z_i}) \cap D(z_j,\delta'_{z_j})$, for any $z_1, z_2 \in J_2$
Therefore F' can be continuously extended in J_2

Next we prove statement (b):

We have already that $F' = [\phi \circ (\phi^{-1} \circ F)]' = \phi' \circ (\phi^{-1} \circ F) \cdot (\phi^{-1} \circ F)'$
Let $z \in J_2$. Since for all z_n accumulating in J_2, the sequence $F(z_n)$ accumulates in γ, it follows that $\phi^{-1} \circ F(z) \in J_1$. Assuming that $\phi' \neq 0$, for all z in J_1, it follows that $\phi' \circ (\phi^{-1} \circ F)(z) \neq 0$ for all z in J_2.
From the Reflection Principle, the function $h = \phi^{-1} \circ F$ can be extended in an neighbourhood of J_2 and the extension is injective on it; thus $(\phi^{-1} \circ F)' \neq 0$, for all z in J_2.Therefore $F'(z) \neq 0$ for all $z \in J_2$. This completes the proof.

□

Remark 2.3. Let suppose that there exists z in J_1 such that $\phi'(z) \neq 0$ and not for all points in J_1. Therefore, there exists ζ in J_2 such that $\phi^{-1} \circ F(\zeta) = z$ in J_1 and $F(\zeta) \neq 0$ and reversely.

Lemma 2.4. *Let $W \subseteq \mathbb{C}$ and $\Omega \subseteq \mathbb{C}$ open sets. Let $\phi : W \to \mathbb{C}$ and $h : \Omega \to W$ holomorphic functions. We assume that P is a finite linear combination of products of the functions $\phi^{(n)} \circ h, \phi^{(n-1)} \circ h, ..., \phi \circ h$ and $h^{(n)}, ..., h$, for some n in $\mathbb{N}$. Then P is holomorphic on Ω and its derivative is a polynomial of the functions $\phi^{(n)} \circ h, \phi^{(n-1)} \circ h, ..., \phi \circ h$ and $h^{(n)}, ..., h$ and $\phi^{(n+1)} \circ h, h^{(n+1)}$*

Proof. We will use induction to prove this lemma. We start from

$$P(z) = (\phi \circ h)^{k_1} h^{k_2}(z)$$

, then

$$P'(z) = k_1(\phi' \circ h)h'h^{k_2} + k_2(\phi \circ h^{k_1})h^{k_2-1}h'$$

For the inductive step we assume

$$P = c[\phi^{(n-1)} \circ h^{k_{n^2}} \dots \phi \circ h^{k_{n+1}} h^{(n)^{k_n}} \dots h^{k_1}], c \in \mathbb{C}$$

Without loss of the generality, we also assume that $k_{2n} = \dots = k_1 = c = 1$, hence,

$$\begin{gathered} P' = (\phi^{(n)} \circ h)h'[\phi^{(n-1)} \circ h) \dots (\phi^{-1} \circ F)] \\ +(\phi^{(n-1)} \circ h)^2 h'(\phi^{(n-2)} \circ h \dots h) \\ + \dots + [\phi' \circ h]^{(n+1)}([\phi^{(n)} \circ h] \dots h) \\ +h^{(n+1)}([\phi^{(n)} \circ h]) + \dots + h([\phi^{(n)} \circ h] \dots h') \end{gathered}$$

Therefore P' is a polynomial of $(\phi^{(n)} \circ h)^{(k)}, h^{(k)}, k = 0, ..., n+1$ □

Corollary 2.4.1. *According to assumptions and notation of Theorem 2.2, if ϕ is an analytic function in W_1 and continuous on J_1 and we also assume that $\phi^{(k)}$ can be continuously extended on J_1 for $k = 0, 1, ..., p$, where p is a natural number, then $F^{(k)}$ can be continuously extended on $W_2 \cup J_2$ for $k = 0, 1, ..., p$.*

This holds because $F^{(k)}$ is a polynomial of functions that can be continuously extended on J_2. This follows combining Theorem 2.2 with lemma 2.4. (for more details see [3])

The Theorem 2.2 implies, also, the following corollary.

Corollary 2.4.2. *We suppose that $D(0, r, 1), 0 < r < 1$ is an open annulus and $\phi : \overline{D(0, r, 1)} \to \mathbb{C}$ is continuous, injective on $\overline{D(0, r, 1)}$ and holomorphic on $D(0, r, 1)$. We also assume that ϕ' has a continuous extension on $\overline{D(0, r, 1)}$. Furthermore, we set $\gamma(e^{i\theta}) := \phi(e^{i\theta})$ and W is the interior of the*

Jordan curve γ.
We, finally, assume that $\phi(D(0,r,1)) \subseteq W$ and the function

$$F : D(0,1) \to W$$

is a conformal mapping of $D := D(0,1)$ onto W. It is known that according to the Ozgood-Caratheodory theorem ,F is exteded to a homeomorphism $F : \overline{D(0,1)} \to \overline{W}$. From the Theorem 2.2, it follows that

1. *The function F' extends continuously on $\overline{D}$*
2. *If $\phi'(z) \neq 0$, for some z with $|z| = 1$ then*
   $F'(\zeta) \neq 0$, for ζ such that $(\phi^{-1} \circ F)(\zeta) = z$

More generally, if a conformal collar of a Jordan curve has some nice properties, then the same holds for any other conformal collar of the same curve from the same side.

References

[1] Ahlfors, Complex analysis,Second edition ,McGraw-Hill,New York,1966

[2] Steven R.Bell and Steven G Krantz, Smoothness to the boundary of conformal maps, Rocky Mountain Journal of mathematics, Volume 17,1987, Number 1 ,23-40

[3] E.Bolkas,V.Nestoridis,C. Panagiotis and M.PapadimitrakisOne sided extendability and p-continuous analytic capacities,arxiv : 1606.05443

[4] Gauthier,P.M., Nestoridis V., Conformal extensions of functions defined on arbitrary subsets of Riemann Surfaces. Arch. Math. (Basel) 104 (2015)no 1 , 61-67.

[5] Georgakopoulos N. ,Extensions of the Laurent Decomposiotion and the spaces $A^p(\Omega)$, arXiv:1605.08289

[6] Koosis P. ,An introduction to H_p spaces, Cambridge University Press, 1998

[7] V.Mastrantonis, Relations of the spaces $A^p(\Omega)$ and $C^p(\partial\Omega)$, arXiv:1611.02971

National and Kapodistrian University of Athens,
Department of Mathematics
15784
Panepistemiopolis
Athens GREECE
e-mail: lvda20@hotmail.com
e-mail: vnestor@math.uoa.gr